

## EFFECT OF DIFFERENT DOSAGES OF BACTERIA AFTER MULTIPLICATION ON SEAWEED IN THE CONTROL OF ROOT ROT OF SUNFLOWER AND MASHBEAN

Shahnaz Dawar<sup>1</sup>, Sadia Arjumand<sup>1</sup>, M. Javed Zaki<sup>1</sup> and Muhammad Abid<sup>2</sup>

<sup>1</sup>Department of Botany, University of Karachi, Karachi-75270, Pakistan

<sup>2</sup>Department of Botany, Federal Urdu University of Arts, Science and Technology, Gulshahn-e-Iqbal, Karachi, Pakistan

---

### ABSTRACT

Bacteria viz., *Rhizobium meliloti*, *Pseudomonas aeruginosa* and *Bacillus subtilis* were multiplied on sea weed like *Cystoclanium purpuraeum*. After 15 days of multiplication soil was amended with different dosage of inoculum. There was a significant enhancement of plant growth and significantly control the infection of root infecting fungi viz., *Macrophomina phaseolina*, *Rhizoctonia solani* and *Fusarium* spp on mash bean where soil was amended with *P. aeruginosa* inoculum on *Cystoclanium purpuraeum* used @ 1% w/w whereas *B. subtilis* inoculum on *C. purpureum* used @ 1% w/w showed greater plant growth and significant reduction of root infecting fungi on sunflower and mashbean.

**Key words:** Sea weeds, root rot fungi, bacteria, sunflower and mash bean.

---

### INTRODUCTION

Many plant growth promoting rhizobacteria (PGPRS) have a beneficial effect on plants including biological control of soil borne pathogens, induced systemic resistance to plant pathogens, phytoharmonic productions and improvement of nutrition and water uptake of plants (Seuk Bae *et al.*, 2000). Of the *Pseudomonas* species, *P. aeruginosa* has been found as an effective biocontrol agent of root rot pathogens (Izhar *et al.*, 1995) similarly *Rhizobium* spp., the root nodulating bacteria, are also known to control soil borne root infecting fungi (Zaki and Gaffar, 1987; Siddiqui *et al.*, 1998). Seaweeds provide better plant growth and received the attention of plant scientists through out the world (Atzmon *et al.*, 1994, Staden *et al.*, 1995). Since it contain greater potash and nitrogen as compared to farm yard manure (Chapman and Chapman, 1980).

Sea weeds also known to stimulate the growth of vegetables, fruits and other crops (Bhenden., 1991; Crouch *et al.*, 1994; Washington *et al.*, 1999). *Stoechospermum marginatum* and *Sargassum tenerrimum* with or without Rhizobia significantly reduced root knot nematode (*Meloidogyne javanica*) and root infecting fungi (Ehtesham-ul-Haque *et al.*, 1996).

Plant diseases causing organisms produces serious losses to crop plants and adversely affects the agricultural economy of a country (Hafeez, 1986). The primary diseases threatening crop production are due to fungi, actinomycetes, bacteria and nematodes. These are ubiquitous soil borne plant pathogens, which infect roots of plant, resulting in the death of plants. Since damage to plants by soil borne pathogens results from below ground infection, losses to crop plants from such diseases are underestimated and generally go unnoticed (Baker and Cook, 1974). The soil borne root-infecting fungus, *Macrophomina phaseolina* is reported to produces charcoal rot over 500 species of plants (Sinclair, 1982). *Rhizoctonia solani* exists as active mycelium in the soil, attacks over 2000 species of plants (Parmeter, 1970) and *Fusarium* species (Booth, 1971) are known to attack a wide range of host plants in different parts of the world.

Experiments were therefore carried out to study the effect of different dosage of bacteria after multiplication on sea weed in the control of root rot of sunflower (*Helianthus annuus* L.) and mash bean (*Vigna mungo* (L.) Hepper).

### MATERIALS AND METHODS

Seaweed, *C. purpuraeum*, collected from Buleji, Karachi, was washed with tap water and dried under shade and powdered in an electric grinder and was kept at room temperature. Cultures of *Rhizobium meliloti* (R5), *Pseudomonas aeruginosa* (P-58), *Bacillus subtilis* (B-35) were obtained from culture collection of Department of Botany, University of Karachi.

Soil used for the experiment was obtained from the experimental plots of the Department of Botany, University of Karachi and sieved through 2mm sieve to discard particles. The soil used was sandy loam (Sand, Silt, Clay; 70, 19, 11%), pH range from 7.5 - 8.1 with moisture holding capacity (MHC) of 24.04 % (Keen and Raczkowski, 1922), total nitrogen 1.5 % (Mackenzie and Wallace, 1954), total organic matter 2.4 %. Soil had natural infestation of 1-3

sclerotia of *M. phaseolina* as found by wet sieving dilution technique (Sheikh and Ghaffar, 1975), 5-10% colonization of *R. solani* on sorghum seeds used as baits (Wilhelm, 1955) and 3000cfu of *Fusarium* spp., as assessed by soil dilution technique (Nash and Snyder, 1962).

Seaweed (*C.purpuraeum*) was taken in flasks, moistened with water slightly and inoculated with cell suspension of bacteria viz., *R.meliloti* (R5), *P.aeruginosa* (P-58), and *B.subtilis* (B-35) and kept at room temperature ( $25 \pm 2$  °C). After 2 weeks of growth of bacteria, soil was amended with inoculum of *R. meliloti* ( $5.4 \times 10^9$  cells/ml) *P. aeruginosa* ( $7.7 \times 10^9$  cells/ml) and *B.subtilis* ( $4.2 \times 10^9$  cells/ml) @ 0.1%, 0.5%, 1% and transferred in 8cm diam., plastic pots each containing 300gm soil. After 2 weeks of the decomposition of the organic matter 8 seeds of mash bean and sunflower were sown in each pot. Pot with out bacteria and sea weeds served as control. There were three replicates of each treatment pots were randomized on a green house bench. After 30 days, roots of sunflower and mash bean were washed in a running tap water, surface sterilized in 1% Ca (OCl)<sub>2</sub>, and then five 1 cm long root pieces were transferred onto PDA plates containing penicillin @100,000/litre and streptomycin @ 20mg/l. Petri dishes were incubated for 5 days at room temperature to confirm infection of root infecting fungi.

Data were analyzed and subjected to analysis of variance (ANOVA) following the procedure as given by Gomez and Gomez (1984).

## RESULTS AND DISCUSSION

In the present study effect of different dosage of bacteria after multiplication on sea weed in the control of root rot of sunflower (*H. annus* L.) and mash bean (*V. mungo* (L.) Hepper). Greater plant length was observed in both mash bean and sunflower where *P. aeruginosa* and *B. subtilis* inoculum on *C. purpuraeum* used @ 1% w/w ( $p < 0.001$ ). Similarly there was a significant increased in shoot weight, root weight of mash bean and sunflower ( $p < 0.05$ ). Maximum shoot and root weight were observed in sunflower where *P. aeruginosa* inoculum on *C. purpuraeum* used @ 0.5 and 1% w/w. Different dosage of bacteria after multiplication on sea weed were more effective in the suppression of *Fusarium* spp., infection on mash bean ( $p < 0.05$ ) and sunflower ( $p < 0.01$ ) (Table 1). Bacterial inoculum used @ 1% w/w showed complete suppression of *R. solani* infection on mash bean and sunflower ( $p < 0.01$ ). Similarly infection of *M. phaseolina* was significantly reduced when *P. aeruginosa* and *B. subtilis* inoculum on *C. purpuraeum* used @0.5% w/w in mash bean and sunflower ( $p < 0.01$ ). (Table 1).

Table. 1 Effect of different dosage of biocontrol bacteria multiplied on *Cystoclonium purpuraeum* on growth parameters and infection % of mash bean and sunflower.

### Mash bean

Treatments	Growth parameters				Infection %		
	Shoot length (cm)	Shoot weight (gm)	Root length (cm)	Root weight (gm)	<i>Fusarium</i> spp	<i>Macrophomina phaseolina</i>	<i>Rhizoctonia solani</i>
A	17.8	0.8	7.1	0.38	100	100	100
B	21.3	0.67	6.4	0.446	100	100	100
C	19.7	0.73	5.96	0.58	100	83.33	100
D	19.6	0.66	4.16	0.58	83.33	66.66	83.33
E	20.8	0.96	5.96	0.5	833.33	83.33	83.33
F	20.6	0.86	13.6	0.49	66.66	66.66	83.33
G	24.6	1	4.3	3.4	50	100	66.66
H	19.8	0.67	4.5	0.39	66.66	83.33	83.33
I	14.2	0.68	3.66	0.35	66.66	83.33	66.66
J	21	0.67	6.2	0.37	66.66	83.33	50
K	17.7	0.86	4.86	0.34	83.33	66.66	66.66
L	22.3	0.84	4.76	1.82	83.33	50	33.33
M	28.5	1.27	7.1	0.71	33.33	50	16.66
Lsd0.05=	4.28	0.32	2.83	1.71	39.52	38.35	39.14

Sun flower

Table 1 cont'd...

Treatments	Growth parameters				Infection %		
	Shoot length (cm)	Shoot weight (gm)	Root length (cm)	Root weight (gm)	<i>Fusarium</i> spp	<i>Macrophomina phaseolina</i>	<i>Rhizoctonia solani</i>
A	18.7	5.63	0.53	0.85	100	100	100
B	17.2	3.86	0.54	1.23	100	83.33	100
C	21.2	4.96	0.43	0.62	100	83.33	100
D	22.7	5.86	0.48	1.03	83.33	83.33	83.33
E	21.1	5.76	0.52	0.88	83.33	100	83.33
F	20.8	4.44	0.39	0.86	83.33	66.66	50
G	23.5	7.7	0.36	1.2	66.66	83.33	33.33
H	20.4	6.33	0.45	0.71	100	83.33	83.33
I	21.7	4.83	0.40	1.26	83.33	83.33	83.33
J	28.2	9.73	2.18	1.64	66.66	0	16.66
K	18.6	5.4	0.57	42.7	83.33	83.33	83.33
L	20.2	9.06	0.5	1.85	83.33	66.66	66.66
M	30.2	7.63	0.883	1.75	66.66	50	83.33
Lsd0.05=	3.73	3.21	0.31	0.77	34.39	39.71	37.35

A=Control, B=*C. purpuraeum* @ 0.1% w/w, C=*C.purpuraeum* @0.5% w/w, D=*C.purpuraeum* @ 1% w/w, E=*R.meliloti* inoculum @ 0.1% w/w, F=*R.meliloti* inoculum @ 0.5% w/w, G=*R.meliloti* inoculum @1% w/w, H=*P.aeruginosa* inoculum @ 0.1% w/w, I=*P.aeruginosa* inoculum @ 0.5% w/w, J=*P.aeruginosa* inoculum @ 1% w/w, K=*B.subtilis* inoculum @ 0.1% w/w, L=*B.subtilis* inoculum @ 0.5 % w/w., M=*B.subtilis* inoculum @ 1% w/w.

Microorganisms that can grow in the rhizosphere are ideal for use as biocontrol agents since rhizosphere provides front line defense for roots against attack by pathogens (Weller, 1998). Present study on the efficacy of biocontrol bacteria after multiplication on sea weed substrate showed correlation between reduction in infection and population of microbial antagonists. Plant growth promoting rhizobacteria that colonize roots have been reported to improve plant growth either through direct stimulation of the plant by producing growth regulators or by suppression of pathogens (Raaijmaker *et al.*, 2002 ; Weller *et al.*, 2002). The production of certain antibiotics (Leavy *et al.*, 1992) and siderophores (Buysens *et al.*, 1996) by *P. aeruginosa* has been regarded as one of the mechanism involved in antagonism. Siddiqui *et el* (2000) and Siddiqui and Ehtesham-ul-Haque (2001) reported that species of *Pseudomonas* are to be antagonistic to soil borne plant pathogens. Raaijmaker and Weller, (1998) reported role of 2, 4 diacetylpheyloroglucinol an antifungal metabolite from species of fluorescent *Pseudomonas* in plant root disease suppression. Species of *Pseudomonas* are also reported to induce systematic resistance in cucumber against *Pythium aphanidermatum* (Zhou and Paulitz, 1994). Present results would suggest that inoculum of *P. aeruginosa* and *B. subtilis* on *C. purpuraeum* (@ 1% w/w) have greater potential in controlling the soil borne root infecting fungi on mash bean and sunflower.

## ACKNOWLEDGMENT

This work is carried out under the grant of Dean Faculty of Science which is sincerely acknowledged.

## REFERENCES

- Atzmon, N., J.V. Staden and S.J. Van (1994). The effect of sea weed concentration on the growth of *Pinus pinea* seedlings. *New Forests*, 8l: 279-288.
- Baker K.F. and R.J.Cook (1974). *Biological control of plant pathogen*. W.H. Freeman press, San Francisco.
- Booth, C. (1971). *The genus Fusarium*. Common Wealth Mycological Institute, Kew, Surrey England. pp 237.
- Blunden, G. (1991). *Agricultural uses of sea weeds and sea weed extracts. Sea weeds resource in Europ. Uses and potential*. John Willey & Sons, Chichester, pp. 65-81.
- Buysens, S., K. Heungens, J. Poppe and M. Hofte (1996), Involvement of pyochelin and pyoverdine in suppression of *Pythium*-induced damping-off of tomato by *Pseudomonas aeruginosa* TNSK2. *Appl. Environ. Microbiol.*, 62: 865-871.

- Crouch, I. J., J. Van Staden, J. Van. and J. Van Staden (1994) Commercial seaweed products as biostimulants in horticulture. *J. Home Consum. Hort.*, 1: 19-76.
- Chapman, V.J. and D.J. Chapman (1980). *Sea weed and their uses*. 3<sup>rd</sup> ed. Chapman and Hall, New York. pp.334.
- Ehteshamul-Haque, S., M. Abid, V. Sultana, J. Ara and A. Ghaffar (1996). Use of organic amendments on the efficacy of biocontrol agents in the control of root rot and root knot disease complex of okra. *Nematol. medit.*, 24: 13-16.
- Gomez, K.A and A.A.Gomez (1984). *Statistical procedures for agriculture research* 2<sup>nd</sup> ed. Wiley, New York pp.680.
- Hafeez, A. (1986). *Plant diseases*. Pakistan Agricultural Research Centre (PARC), Islamabad. pp 552.
- Izhar, I.S. Ehteshamul-Haque, M.Javed and A.Ghaffar (1995). Efficacy of *Pseudomonas aeruginosa* and *Bradyrhizobium* sp., in the control of root disease in chick pea. *Pak.J.Bot.*, 27: 451-455.
- Keen, B.A. and H. Rackowski (1922). The relation between clay content and certain physical properties of soil. *J. Agric. Sci.*, 11:441- 449.
- Leavy, E., Z. Eyal, I. Chet and A. Hochman (1992). Resistance mechanisms of *Septoria tritici* to antifungal products of *Pseudomonas*. *Physiol. Mol. Pl. Pathol.*, 40: 163-171.
- Mackenzie, H.A. and H.S.Wallace (1954). The Kjeldahl determination of nitrogen: A critical study of digestion conditions, temperature, Catalyst and oxidizing agents. *Aust.J.chem.*, 7:55-70.
- Nash, S.M. and W.C.Snyder (1962). Quantitative estimation by plate counts of propagules of the bean root rot *Fusarium* in field soils. *Phytopathology*, 52:567-572.
- Parmeter, J.R. (1970). *Rhizoctonia solani biology and pathology*. Univ. of California Press, Berkeley Los Angeles and London. pp. 255.
- Raaijmakers, J. M. and D. M. Weller (1998). Natural plant protection by 2, 4-diacetylphloroglucinol producing *Pseudomonas* spp. in takeall decline soils. *Mol. Plant-Microbe Interact.*, 11: 144-152.
- Raaijmakers, J. M., M. Vlami and J. T. de Souza (2002). Antibiotic production by bacterial biocontrol agents. *Antonie van Leeuwenhoek*, 81: 537-547.
- Sheikh, A.H. and A. Ghaffar (1975). Population study of sclerotia of *Macrophomina phaseolina* in cotton field. *Pak. J. Bot.*, 7:13-17.
- Seuk. Bae, Yeoung, OK-Hee Choi, Kyung-Soek park, Sang-Bum Lee and Choong-Hoe Kim (2000). *A useful method for functional analysis of plant growth promoting rhizobacteria in the development of cucumber root system*. Plant pathology Division, National Institute of Agricultural Science and Technology, Suwon 441-707, Korea.
- Siddiqui, I.A., S.Ehteshamul-Haque and A.Ghaffar (1998). Effect of Fungicides on the efficacy of *Rhizobium meliloti* and *Bradyrhizobium* sp., in the control of root infecting fungi on chick pea. *Pak.J.Bot.*, 30:69-74.
- Siddiqui, I. A., S. A. Qureshi, V. Sultana, S. Ehteshamul-Haque and A. Ghaffar (2000) Biological control of root rot root knot disease complex of tomato. *Plant Soil*, 227: 163-169.
- Siddiqui, I. A. and S. Ehteshamul-Haque (2001). Multiplication of *Pseudomonas aeruginosa* on talc-based formulation for use in the control of root rot and root knot disease in tomato. *Trop. Sci.*, 41: 163-167.
- Sinclair, J.B. (1982). *Compendium of soyabean diseases*, 2<sup>nd</sup> ed., American Phytopathological Society. pp.104.
- Staden, J.V., R.P. Beckett, M.J. Rijkberg and J.S. Van (1995). Effect of sea weed concentration on the growth of seedlings of three species of *Eucalyptus*. *South-African. J. Bot.*, 61: 169-172.
- Washington, W. S., S. Engleitner, G. Boontjes and N. Shanmuganathan (1999). Effect of fungicides, seaweed extracts, tea tree oil and fungal agents on fruit rot and yield in strawberry. *Aust. J. Exp. Agric.*, 39: 487-494.
- Weller, D.M. (1998). The role of antibiosis in the biological control of plant disease. *Ann. Rev. Phytopath.*, 26: 75-91.
- Weller, D. M., J. M. Raaijmakers, B. B. M. Gardener and L. S. Thomashow (2002). Microbial population responsible for specific soil suppressiveness to plant pathogens. *Annu. Rev. Phytopathol.*, 40: 309-348.
- Wilhelm. S. (1955). Longevity of the *Verticillium* wilt fungus in the laboratory and field *Phytopathology*, 45:180-181.
- Zaki, M.J and A.Ghaffar (1987). Effect of *Rhizobium* spp., on *Macrophomina phaseolina*. *Pak.J.Sci. Ind. Res.*, 30: 305-306.
- Zhou, T., and T. C. Paulitz (1994). Induced resistance in the biocontrol of *Pythium aphanidermatum* by *Pseudomonas* spp., on cucumber. *J. Phytopath.*, 142: 51-63.

(Accepted for publication October 2006)