

Prevalence of *Helicobacter pylori* infection causing serious gastric problems in individuals consuming spicy food in district Khairpur, Pakistan

Nisar Abbas Shar, Rafique Abbas Shar, Gul Afshan Soomro, Ghulam Abbas Shar, Muhammad Bachal Korai, Shaista Khan

Department of Medicine, Ghulam Mohammad Mahar, Medical College Hospital
Shah Abdul Latif University, Khairpur, Sindh, Pakistan

Objective: To investigate the prevalence of *H. pylori* in spicy food consuming individuals of district Khairpur, Pakistan.

Methodology: In this cross-sectional study, total of 238 patients were included and samples of the blood were taken. The people were interrogated by utilizing the planned pre-tested questionnaire. Immunochromatographic technique (ICT) kit method was used for serum antibodies. Immunoglobulin acute & chronic (IgM/IgG) test was used for antibodies in samples. Data were analyzed on SPSS version 17.

Results: The rate was allied with sort of consuming spicy food in adult with $p = 0.00001$. *H. pylori*

infection exhibited substantial association with spicy food intake. In 238 individuals consuming spicy food, 158 were positive for *H. pylori* IgM/IgG antibodies. Overall, *H. pylori* was seen in 66.38% cases.

Conclusion: The incidence of *H. pylori* is considerably high in individuals with intake of spicy foods ($p < 0.0001$) in our district. However, further studies are suggested to determine the other risk factors for acquisition of *H. pylori* infection in this area.

Keywords: *Helicobacter pylori*, immunochromatography technique, IgG antibodies.

INTRODUCTION

H. pylori is a gram-negative bacterium, generally present in human stomach and causes infection.^{1,2} In 1980's Warren and Marshall from Perth, Australia reported culture a bacterium that was *Campylobacter pyloridis*. *Helicobacter* Genus was formed in 1989 and was given the title as *H. pylori*.³ Its size is 0.5 – 1.0 μm width and 2.5 – 5.0 μm length and has 4 – 6 flagella.⁴ Inside stomach, most of the *H. pylori* is detectable within the gastric mucosa; a few, however, are adherent to intestinal epithelial mucosa.⁵ Transmission of *H. pylori* from person to person by the oral-oral or fecal-oral route is most likely.⁶ Bacteria have been isolated from stool, saliva, and dental plaque of infected people.⁷ The incidence of *H. pylori* and related infections has been extremely unpredictable globally.⁸

In developing countries of Asia including Pakistan, India, Thailand and Bangladesh the incidence of *H. pylori* infection is greater while in developed countries of Asia as China and Japan it was shown in the early era.⁹ Humans have an inhospitable stomach and usually fasting conditions do not support any bacteria except few lactobacilli and *H. pylori*.¹⁰ The *H. pylori* is highly adaptive which allow it to colonize.¹¹

Spicy food consumers adults have symptoms of dyspepsia that are greatly linked to this infection.¹² The rate among adult are higher than 80% in various

developing countries versus 25% in industrial nations.¹³ *H. pylori* infection causes chronic gastritis, peptic ulcer and gastric cancer.¹⁴ Epidemiological prevalence of infection of *H. pylori* is not available in Khairpur. Therefore, the need of the hour was to investigate the gastric problems to take timely efforts for controlling the spread of such disease.

METHODOLOGY

In this cross-sectional study, 238 patients were included and samples of the blood were taken. The people were interrogated by utilizing the planned pre-tested questionnaire and blood samples were collected using standard protocol. Immunochromatographic technique (ICT) kit method IgM enzyme-linked immunosorbent assay (ELISA kit, Cat # EIA-2111, Tests: 96 wells, DRG Instruments GmbH, Germany) was used for analysis of samples.

An authorized official permission letter was acquired from the Head of Medicine Department of KMC Hospital Khairpur and Pir Syed Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat (GIMS), Khairpur.

Statistical Analysis: Data were analyzed on SPSS version 17. Chi square, p-value, and Pearson Chi-square tests were statistically conducted.

RESULTS

During the survey study, 334 respondent patients were interviewed using pre organized questionnaire to collect the information and they answered several questions regarding the symptoms suffering from. The obtained results were compiled and analyzed. Total number of respondent patients during survey was 334. Out of 344 total respondent patients, 95.2% (318/334) had nausea, 94.6% (316/334) had vomiting, 94.3% (315/334) heartburn and abnormal bloating, 84.7% (283/334) poor appetite, 82% (274/334) weight loss. We found that 84.7% (283/334) individuals consumed spicy food (Table 1). Incidence of *H. pylori* in current study was higher significantly 84.7% in those whose dietary habits were spicy food than those 15.3% whose dietary habits were non-spicy food.

Out of 334, 238 respondents were agreed for consent to perform their tests for prevalence of *H. pylori* and 96 were disagreeing. IgM/IgG test ICT kit test for *H. pylori* in spicy food consuming patients showed positive results as 66.38% (158/238) and negative as 33.61% (80/238) as shown in Table 2.

Table 1: Respondents having spicy food and symptoms.

Symptoms	Yes	No	Percentage	
			Yes	No
Having Spicy Food	283	51	84.7	15.3
Nausea	318	16	95.2	4.8
Vomiting	316	18	94.6	5.4
Heart Burn and Abnormal	315	19	94.3	5.7
Poor Appetite	283	51	84.7	15.3
Weight loss	274	60	82	18
Previous History	116	218	34.7	65.3

Table 2: Result of IgM/IgG Test for *H. pylori* in serum.

Positive		Negative		Total
Number	%	Number	%	
158	66.38%	80	33.61%	238

Statistically analysis of data compared for validation of our results and Chi-square results were less than 0.014 (Table 3). Table 3 also shows the *H. pylori* associated with spicy food factors in which p-value was 0.000 for spicy food consumption.

Table 3: Validation and Significance (p-value) of *H. pylori* results by SPSS.

Method	Significance
Pearson Chi-Square	0.014
Significance (p-value)	
Spicy Intake	Significance (p-value)
	0.00001

DISCUSSION

There are various symptoms like nausea, vomiting, heartburn, poor appetite etc., by which the prevalence of *Helicobacter pylori* can be predicted. The study in Kassala State, East of Sudan, revealed that among 431 schoolchildren the most common signs linked to *H. pylori* infections was nausea in 25.5%.¹⁵ Another study revealed that nausea was considerably allied with the *H. pylori*.¹⁶ A greater correlation with active *H. pylori* infection was found in 93.3% for alcohol drinking and 80% for non-vegetarian food consumption.¹⁷ The results of our study showed that many respondents had nausea, vomiting, heartburn and abnormal bloating, poor appetite and weight loss (Table 1).

Infection with *H. pylori* may also be linked with diet and feeding practices.¹⁸ Eating kipper and fried food was positively associated with *H. pylori* infection. Furthermore, salty food itself may be a source of *H. pylori*.¹⁹ The synergistic amplifying influence of infection with *H. pylori* and consumption of salty foods was noted in a case control study in Korea.²⁰ In a previous study, individuals who consumed marinated foods infrequently than on any second day were the most highly infected (66.77%).¹⁹ In our study, 84.7% individuals consumed spicy food (Table 1).

A previous study on *H. pylori* by an enzyme-linked immunoassay revealed that 0.82 (0.74 – 0.91) – fold and 0.88 (0.79 – 0.98) – fold greater risk of *H. pylori* seropositivity in smokers and drinker than those who had never consumed them.²¹ Another study showed total IgG antibody seropositivity was 79.3 percent, rank correlated inversely with infection and non cigarette smoking and drinking alcohol were associated with infection.²² A cross-sectional study of 634 men randomly selected in Japan and the 474 men were positive for the IgG antibody against the bacterium.²³

H. pylori infection in dyspeptic patients seems to be connected with gastritis in a study from Islamabad showed the 94.6% patients with chronic gastritis.²⁴ Statistically analysis of our data in SPSS compared the validation of our results with chi square and Pearson Chi-square result less than 0.014. The *H. pylori*

associated with spicy food factor in which *p*-value was found to be 0.00001 for spicy food consumption (Table 3).

CONCLUSION

The rate of *H. pylori* infection was relatively high in males (68%) than females (32%). The blood serology test results of individuals consuming spicy food were positive for *H. pylori* IgM/IgG in district Khairpur. Better understanding the prevailing pattern will help in prioritizing public health efforts to effectively manage *H. pylori* infection associated complications.

Author Contributions:

Conception and design: Nisar Abbas Shar, Ghulam Abbas Shar.
Collection and assembly of data: Nisar Abbas Shar, Ghulam Abbas Shar.
Analysis and interpretation of data: Rafique Abbas Shar, Gul Afshan Soomro, Muhammad Bachal Korai.

Drafting of the article: Nisar Abbas Shar, Rafique Abbas Shar, Shaista Khan.

Critical revision of article for important intellectual content: Nisar Abbas Shar.

Statistical expertise: Gul Afshan Soomro, Muhammad Bachal Korai.

Final approval and guarantor of the article: Ghulam Abbas Shar.

Corresponding author email: Ghulam Abbas: gabbas.shar@salu.edu.pk

Conflict of Interest: None declared.

Rec. Date: Jun3, 2021 Revision Rec. Date: Nov9, 2021 Accept Date: Dec 3, 2021.

REFERENCES

- Hooi JK, Lai WY, Ng WK, Suen MM, Underwood FE, Tanyingoh D, et al. Global prevalence of *Helicobacter pylori* infection: systematic review and meta-analysis. *Gastroenterology*. 2017; 153: 420-9.
- Driscoll LJ, Brown HE, Harris RB, Oren E. Population knowledge, attitude, and practice regarding *Helicobacter pylori* transmission and outcomes. *Front Public Health*. 2017; 5: 144-9.
- Ali MM. Molecular Detection of Metronidazole Resistant Genes (RdxA and FrxA) of *Helicobacter pylori* Isolated from Stool Specimen at Alinjaz Diagnostic Center in Khartoum State (Doctoral dissertation, Sudan University of Science & Technology), 2019.
- Viji S. A Study on Diagnosis of *Helicobacter Pylori* Infection by Culture and Molecular Methods from Gastric Biopsy Specimens and Serological Assays in Patients with Peptic Ulcer Disease (Doctoral dissertation, Madras Medical College, Chennai), 2019.
- Ahmed F, Ahmed J, Ahmed H, Hassan SS, Iqbal J, Sharif A. Advances in research progress of *H. pylori*. *J Pharm Res Int*. 2018; 8: 1-8.
- Nouh, F. The affect of salty food intake on the risk of *Helicobacter pylori* infection among patients attending Benghazi medical center (A retrospective cohort study). *Sch J App Med Sci*. 2021; 8: 1274-82.
- Park SY, Hwang BO, Lim M, Ok SH, Lee SK, Chun KS, et al. Oral-Gut micro biome axis in gastrointestinal disease and cancer. *Cancers*. 2021; 13: 2124-43.
- Muhammad JS, Zaidi SF, Sugiyama T. Epidemiological ins and outs of *helicobacter pylori*. *J Pak Med Assoc*. 2012; 62: 955-9.
- Zaidi SF. *Helicobacter pylori* associated Asian enigma: Does diet deserve distinction? *World J Gastrointest Oncol*. 2016; 8: 341-50.
- Kivi M. Aspects of *Helicobacter pylori* transmission. Mikrobiologiskt och Tumörbiologiskt Centrum (MTC)/Microbiology and Tumor Biology Center (MTC); Inst för medicinsk epidemiologi och biostatistik/Dept of Medical Epidemiology and Biostatistics; Swedish Institute for Infectious Disease Control (SMI), 2005.
- Burne RA, Chen YY. Bacterial ureases in infectious diseases. *Microb Infect*. 2000; 2: 533-42.
- Miwa H, Kusano M, Arisawa T, Oshima T, Kato M, Joh T, et al. Evidence-based clinical practice guidelines for functional dyspepsia. *J Gastroenterol*. 2015; 50: 125-39.
- Suerbaum S, Michetti P. *Helicobacter pylori* infection. *N Engl J Med*. 2002; 347: 1175-86.
- Iwai N, Okuda T, Oka K, Hara T, Inada Y, Tsuji T, et al. *Helicobacter pylori* eradication increases the serum high density lipoprotein cholesterol level in the infected patients with chronic gastritis: A single-center observational study. *PloS One*. 2019; 14: e0221349.
- Abbas M, Sharif FA, Osman SM, Osman AM, El Sanousi SM, Magzoub M, et al. Prevalence and associated symptoms of *Helicobacter pylori* infection among schoolchildren in Kassala State, East of Sudan. *Interdiscip Perspect Infect Dis*. 2018; 2018: 1-5.
- Correa Silva RG, Machado NC, Carvalho MA, Rodrigues MA. *Helicobacter pylori* infection is high in paediatric nonulcer dyspepsia but not associated with specific gastrointestinal symptoms. *Acta Paediatr*. 2016; 105: e228-31.
- Khan S, Beevi PF, Husain M, Hassan MJ, Jetley S. *Helicobacter pylori* Infection and Associated Risk Factors in Patients with Upper Gastrointestinal Symptoms. *Morb Mortal Wkly Rep*. 2017; 2: 4.
- Tsugane S. Salt, salted food intake, and risk of gastric cancer: epidemiologic evidence. *Cancer Sci*. 2005; 96: 1-6.
- Zhu Y, Zhou X, Wu J, Su J, Zhang G. Risk factors and prevalence of *Helicobacter pylori* infection in persistent high incidence area of gastric carcinoma in Yangzhong city. *Gastroenterol Res Pract*. 2014; 2014: 1-10.
- Lee SA, Kang D, Shim K, Choe J, Hong W, Choi H. Original article effect of diet and *Helicobacter pylori* infection to the risk of early gastric cancer. *J Epidemiol*. 2003; 13: 162-8.
- Ogihara A, Kikuchi S, Hasegawa A, Kurosawa M, Miki K, Kaneko E, et al. Relationship between *Helicobacter pylori* infection and smoking and drinking habits. *J Gastroenterol Hepatol*. 2000; 15: 271-6.
- Kim JY, Kwon O. Garlic intake and cancer risk: an analysis using the Food and Drug Administration's evidence-based review system for the scientific evaluation of health claims. *Am J Clin Nutr*. 2009; 89: 257-64.
- Tsugane S, Tei Y, Takahashi T, Watanabe S, Sugano K. Salty food intake and risk of *Helicobacter pylori* infection. *JPN J Cancer Res*. 1994; 85: 474-8.
- Mehmood K, Awan AA, Muhammad N, Hasan F, Nadir A. *Helicobacter pylori* prevalence and histopathological findings in dyspeptic patients. *J Ayub Med Coll Abbottabad*. 2014; 26: 182-5.