

## EFFECT OF MAGNESIUM ON SUGARCANE IN PIEDMONT PLAIN SOILS OF BANGLADESH

S.C. Saha & S.R. Ghosh

Sugarcane Research and Training Institute, Ishurdi, Pabna, Bangladesh

Field experiments were conducted on the old Himalayan piedmont plain region of non-calcareous brown flood plain soils (Typic Haplaquept) at two sites, such as Kanta Farm of Shetabgong Sugar Mills, Dinajpur, and North Regional Station (NRS) Farm of Sugarcane Research and Training Institute, Thakurgaon, Bangladesh during two cropping seasons 1991-92 and 1992-93 to study the effect of magnesium on sugarcane. Sugarcane tillering progressively increased under different magnesium levels and attained the highest in May, thereafter gradual reduction occurred due to tiller mortality. The crop responded significantly to magnesium application at both the sites. Percentage increase in cane yield in magnesium treated plots over control ranged from 3.7 to 10.9 at Kanta, and 4.4 to 11.7 at NRS, which was more appreciable at 10 kg Mg ha<sup>-1</sup>. The maximum response dose calculated from cubic equation was 9.5 kg Mg ha<sup>-1</sup> for both tiller stages.

Key words: Bangladesh, magnesium effect on sugarcane, piedmont plain soils

### INTRODUCTION

Until recently, the importance of secondary nutrient elements in general, and magnesium in particular, in Bangladesh agriculture had received little research effort. With the introduction of high-yielding varieties, use of high analysis fertilizers, and adoption of multiple cropping practice in potentially developed areas, deficiency of this nutrient has been increasingly reported. Magnesium deficiency is actually more widespread than it is realized. It is because of the fact that there is very inadequate scientific data about the effect of magnesium fertilizers on crops. Magnesium is a constituent of chlorophyll molecule that is essential for photosynthesis and its application causes good growth of plants especially of foliage (Singh *et al.*, 1981).

Magnesium is of considerable significance in acid soils. Magnesium content in acid soils of Bangladesh is low (Anonymous, 1988-89). Crop responses to applied magnesium have been reported either on acid soils (Tajuddin, 1991) or on sandy soils (Aulakh and Pasricha, 1977; Kumar *et al.*, 1981; Singh and Singh, 1990) which have low content of exchangeable magnesium. Increase in sugarcane yield due to applied magnesium has been reported from Queensland, Australia (Shahjahan, 1980), Puerto Rico (Landrau and Samuels, 1986), and Taiwan (Chao and Li, 1989). There is paucity of information on the effect of magnesium on the yield of sugarcane in Bangladesh. In view of this, the present investigation was conducted to study the effect of magnesium on

sugarcane in the old Himalayan piedmont plain soils of Bangladesh.

### MATERIALS AND METHODS

Field experiments were conducted on the old Himalayan piedmont plain region of non-calcareous brown flood plain soils (Typic Haplaquept) at two sites, namely Kanta Farm of Shetabgong Sugar Mills (Kanta), Dinajpur, and North Regional Station (NRS) Farm of Sugarcane Research and Training Institute, Thakurgaon, Bangladesh during two successive cropping seasons 1991-92 and 1992-93. The experiments were set up in randomized complete block design with four replications. The treatments consisted of four levels of magnesium (0, 10, 20, and 30 kg ha<sup>-1</sup>) as commercial grade magnesium oxide (45%). Location specific recommended doses of N, P, K and S were applied equally to all the treatments. The doses for both Kanta and NRS in kg ha<sup>-1</sup> were: 160N, 120 P<sub>2</sub>O<sub>5</sub>, 150 K<sub>2</sub>O and 40 S. Whole amount of phosphorus, sulphur, and magnesium, and one-third each of nitrogen and potassium were applied at the time of transplanting, the rest two-thirds of nitrogen and potassium were applied in two equal splits of 90 days and 150 days after transplanting. Forty-five days old seedlings of sugarcane variety Isd-16 grown in polybags were transplanted in middle of November at both the sites during both the cropping seasons. All intercultural practices were performed as and when necessary. Tillers were recorded in all the treatments in March, May and July at an interval of 60 days, and the crop was harvested in middle of

January at both the sites. Number of millable cane, percentage recoverable sucrose, and yields of cane were recorded at harvest. Recoverable sucrose percentage was determined according to Horne's dry lead method (Meade and Chen, 1977).

Soil samples (0-25 cm) were collected before start of the experiment at both the locations in both the cropping seasons. The samples were air-dried, ground with a wooden pestle and mortar, and passed through a 2 mm sieve. Soil pH was determined in 1:25 soil: distilled water suspension by pH meter; organic carbon by chromic acid wet oxidation method of Walkley and Black; texture by hydrometer method; total nitrogen by modified Kjeldahl method; available phosphorus was determined after extraction with 0.5M NaHCO<sub>3</sub> at pH 8.5 following chlorostannous reduced molybdophosphoric blue colour method; exchangeable potassium was estimated flame photometrically in NH<sub>4</sub>OAc extract, and exchangeable magnesium in NaOAc extract was estimated by titrating with EDTA solution (Jackson, 1973). Available sulphur was determined after extraction with NaOAc by developing turbidity using BaCl<sub>2</sub>, and gum acacia (Chesnin and Yien, 1950). Characteristics of initial soils are given in Table 1.

Data on tillering and yield parameters of sugarcane, from experiments in each cropping season and location, were subjected to statistical analysis and the significance of mean difference was carried out as per LSD (Gomez and Gomez, 1984). The treatment sums of squares were partitioned into linear, quadratic, and cubic response equation (with pooled yield of Kanta and NRS). Cubic ( $Y = a + bx + cx^2 + dx^3$ ) yield response equation indicating cane yield as a function of applied magnesium was computed, and the response function was computed for efficiency by estimating the coefficient of determination as:  $R^2 = \text{sum of squares due to regression} / \text{sum of squares}$ . From the cubic relationship, maximum response dose was worked out as follows:

Maximum response dose'

$$(\text{kg Mg ha}^{-1}) :: 1/3 d \{-c \pm [C^2 - 3bd]^{1/2}\}$$

Where b, c, and d are the estimates of the regression coefficients.

## RESULTS AND DISCUSSION

Magnesium application had significant effect on tillering of sugarcane during all the recorded months both at Kanta and NRS. Monthly count of tillers starting from 120 days after transplanting revealed

peak tiller population in May which declined as tiller mortality set in thereafter (Table 2). The application of magnesium produced more tillers over control with maximum at 10 kg ha<sup>-1</sup> at both the sites. The yield of sugarcane presented in Table 2 clearly indicates significant response to magnesium application in soil at 10 kg ha<sup>-1</sup> at both the sites. There was a decreasing trend in sugarcane yield due to magnesium application at 20 and 30 kg levels compared to 10 kg level. Percentage increase in cane yield in magnesium treated plots over control ranged from 3.7 to 10.9 at Kanta and 4.4 to 11.7 at NRS, which was more appreciable at 10 kg mg ha<sup>-1</sup>. The economic response was maximum at 10 kg magnesium level both at Kanta (7.1 q ha<sup>-1</sup>) and NRS (7.9 q ha<sup>-1</sup>), and then decreased with an increased level of magnesium at 30 kg (0.8 q ha<sup>-1</sup> at Kanta and 1.0 q ha<sup>-1</sup> at NRS). Similar results with applied magnesium in sugarcane in acid soil were reported by Rahman *et al.* (1986) in Bangladesh and Landrau and Samuels (1986) in Puerto Rico. Chao and Li (1989) reported that the application of magnesium increased yields of sugarcane by 10 to 15 % in Taiwan. The highest yield of cane and sugar obtained with application of 10 kg tJg ha<sup>-1</sup> could be due to deficient levels of magnesium in soil needed to support the growth of sugarcane.

The highest net economic benefit was obtained with 10 kg Mg ha<sup>-1</sup> at both Kanta and NRS (Table 3). The yield data (pooled yield of Kanta and NRS) were also fitted to cubic response curve, of which coefficient of determination ( $R^2$ ) was found to be highly significant. The regression equation was  $Y = 65.0 + 2.7x - 0.1215x^2 + 0.00225x^3$  for  $0 \leq x \leq 30$  ( $R^2 = 0.95^{**}$ ) where, Y is the expected yield (t ha<sup>-1</sup>) and x is the level of magnesium (kg ha<sup>-1</sup>). The results indicate that yield response of sugarcane to magnesium fertilization can be adequately described by cubic equation. The computed  $R^2$  value of 0.953 indicates that 95.3% of total variation in the mean yields was explained by estimated cubic regression equation. Sugarcane responded significantly at 10 kg Mg ha<sup>-1</sup>, but magnesium dose that maximises yield calculated from cubic equation was 9.5 kg ha<sup>-1</sup> with a cane yield of 72.1 t ha<sup>-1</sup>.

The application of magnesium had a significant effect on percentage recoverable sucrose at both the sites. The highest percentage recoverable sucrose was produced by an application of 10 kg Mg ha<sup>-1</sup> followed by 20 kg Mg ha<sup>-1</sup>, and 30 kg Mg ha<sup>-1</sup> at both the sites. Thus, it could be concluded that sugarcane grown on the old Himalayan piedmont plain region of

# Effect of magnesium on sugarcane

Table 1. Characteristics of soils of experimental fields

Character	Kanta*	NRS*
pH	5.40	5.20
Sand (%)	71.00	67.00
Silt (%)	18.00	20.00
Clay (%)	11.00	13.00
Soil texture	Sandy loam	Sandy loam
Organic carbon (%)	0.65	1.15
Total N (%)	0.06	0.08
Available P (ppm)	13.00	12.00
Exchangeable K (me/100 g)	0.11	0.08
Available S (ppm)	10.00	11.50
Exchangeable Mg (me/100 g)	0.60	0.50

\* Average of two years.

Table 2. Effect of magnesium on yield parameters of sugarcane on the old Himalayan piedmont plain soils (average of 2 years)

Location	Treatment (kg Mg ha <sup>-1</sup> )	Tiller population (x 10 <sup>3</sup> ha <sup>-1</sup> )			Millable cane (x10 <sup>3</sup> ha <sup>-1</sup> ) <sup>1</sup>	Yield of cane (ton ha <sup>-1</sup> )	Response/ kg Mg (q ha <sup>-1</sup> )	Recoverable sucrose (%)	Yield of sugar (ton ha <sup>-1</sup> )
		March	May	July					
KANTA	0	60.1	140.4	83.1	71.1	65.0	-	9.2	6.0
	10	67.8	172.8	89.6	76.8	72.1	7.1	9.8	7.1
	20	64.7	169.8	87.2	73.5	68.4	1.7	9.5	6.5
	30	64.0	147.8	84.6	72.6	67.4	0.8	9.4	6.3
	CD (P=0.01)	7.6	7.4	4.3	3.4	2.7	-	0.2	0.2
NRS	0	65.1	143.7	86.1	74.0	67.6	-	9.4	6.4
	10	71.3	176.0	93.1	80.0	75.5	7.9	10.0	7.6
	20	67.9	173.3	90.4	77.0	71.7	2.1	9.7	7.0
	30	66.9	150.8	87.5	76.5	70.6	1.0	9.6	6.8
	CD (P=0.01)	2.4	6.2	6.1	2.4	2.1?	-	0.2	0.2

Table 3. Economics of magnesium fertilization on sugarcane production

Location	Mg levels (kg ha <sup>-1</sup> )	Cost of Mg fertilizer (Taka*)	Yield of cane (ton ha <sup>-1</sup> )	Increased cane yield over control (ton ha <sup>-1</sup> )	Net income (Tk ha <sup>-1</sup> )	Net benefit (Tk ha <sup>-1</sup> )
KANTA	0	-	65.0	-	-	-
	10	3000	72.1	10.9	10900	(+)7900
	20	6000	68.4	5.2	5200	(-) 800
	30	9000	67.4	3.7	3700	(-) 5300
	CD					
NRS	0	-	67.6	-	-	-
	10	3000	75.5	11.7	11700	(+)8700
	20	6000	71.7	6.1	6100	(+)100
	30	9000	70.6	4.4	4400	(-)4600
	CD					

\* Local currency: Rupee one = Taka.

(Cane price Taka 1000.00 t<sup>-1</sup> and magnesium Taka 300.00 kg<sup>-1</sup>).

non-calcareous brown flood plain soils (Typic Haplaquept) of Bangladesh, low in magnesium availability, require's to be fertilized with 9.5 kg Mg ha' along with the recommended doses of N, P, K, and S for better yield.

Acknowledgement: The authors express their thanks to Mr. M.A. Gafur, Associate Statistician at the Sugarcane Research & Training Institute, Ishurdi, for his help in analysis of data. The authors are really grateful to Dr. Ranajit Kundu, Associate Cane Entomologist, SRTI for his constructive suggestions to improve the write up of this article.

## REFERENCES

- Anonymous. 1988-89. Bangladesh Agricultural Research Institute, Annual Report, Joydebpur, Bangladesh.
- Aulakh, M.S. and N.S. Pasricha. 1977. Interrelationships between S, Mg, Kin rapeseed. I. Yield, concentration and uptake of sulphur. *Indian J. Agri. Sci.* 47: 577-581.
- Chao, T.T. and K.Y. Li. 1989. Field experiments on the amelioration of lateritic soil: Effects of lime and magnesia application. Rept. Taiwan Sugar Exp. Sta. No. 4,125-146.
- Chesnin, L. and C.H. Yien. 1950. Turbidimetric determination of available sulphates. *Soil Sci. Soc. Am. Proc.* 15: 149-151.
- Gomez, K.A. and A.A. Gomez. 1984. Statistical Procedures for Agricultural Research. A Wiley-Interscience publication. John Wiley and Sons, New York, USA.
- Jackson, M.L. 1973. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi, India.
- Kumar, V., M. Singh and N. Singh. 1981. Response of wheat to magnesium in sandy soils of Hissar. *Pl. Soil*, 59: 3-12.
- Landrau, P. and G. Samuels. 1986. Results of lime and minor element fertilizer research in Puerto Rico. *J. Agri. Univ. Puerto Rico*, 40: 224-234.
- Meade, G.P. and J.C.P. Chen. 1977. Cane Sugar Handbook. A Wiley-Interscience Publication. John Wiley and Sons, New York, USA.
- Rahman, S.B.M.F., A.B.M.M. Rahman, M.A.A. Sarkar and S.C. Saha. 1986. Calcium and magnesium fertilization of sugarcane. *Bangladesh J. Sugarcane*, 8: 64-66.
- Shahjahan, M. 1980. A report on the visit to Australia. Sugarcane Res. Trg. Inst., Ishurdi, Pabna, Bangladesh.
- Singh, R., B.C. Biswas, S.C. Maheswari and S.C. Srivastava. 1981. Sugarcane. Fertilizer Assoc. of India, New Delhi, India.
- Singh, R.P. and V. Singh. 1990. Effect of S, Mg, and K on yield and uptake by linseed. *J. Indian Soc. Soil Sci.* 38: 169-170.
- Tajuddin, E. 1991. Effect of magnesium on groundnut in acid soils of Kerala. *Agri. Res. J. Kerala*, 28: 89-95.