

Comparison of post-operative pain with and without administration of local anesthetic intraperitoneally in laparoscopic cholecystectomy

Nadia Attiq, Aurangzeb Khan, Muhammad Imran, Khalil-ur-Rehman, Naveed Arshad, Attique Sadique

Department of Surgery, Fauji Foundation Hospital, Rawalpindi, Pakistan

Objectives: To compare post-operative pain with and without administration of local anesthetic intraperitoneally in patients undergoing laparoscopic cholecystectomy.

Methodology: This randomized controlled trial was done in surgical unit of Fauji Foundation Hospital, Rawalpindi from July 2017 to Jan 2018. A total of 120 consecutive patients between age of 20 – 60 years of both gender and who were candidates of laparoscopic cholecystectomy were included in the study. They were equally allocated into two treatment groups by random assignment. Patients in group A (IP 0.5% Bupivacaine) underwent laparoscopic cholecystectomy with administration of 20 ml of 0.5% bupivacaine intraperitoneally. Group B (control) patients underwent laparoscopic cholecystectomy without administration of intraperitoneal local anesthetic.

Efficacy ($VAS \leq 4$) in terms of reduced pain after instillation of local anesthetic intraperitoneally was measured at 8 hours after surgery.

Results: The mean age of both groups was 46.80 ± 7.12 and 49.40 ± 6.19 years, respectively. According to ASA classification I & II, 52 (86.7%) were in group A and 8 (13.3%) in group B. Effective pain control at 8 hours after surgery was achieved in 42 (70%) patients in group A as compared to 12 (20%) in group B ($p = 0.001$).

Conclusions: Intraperitoneal administration of 0.5% Bupivacaine after laparoscopic cholecystectomy was found to be an effective option for post-operative pain management at 8 hours after surgery.

Keywords: Cholecystectomy, Laparoscopic injections, intraperitoneal, postoperative pain.

INTRODUCTION

Cholecystectomy is the most commonly performed procedure on biliary tract and second most commonly performed surgical procedure.¹ Better pain management and reduced hospital stay has made laparoscopic cholecystectomy (LC) the gold standard procedure for symptomatic gall stones.^{2,3} Despite the common belief, LC is still associated with a considerable post-operative pain. Compared with conventional open cholecystectomy where the pain is parietal in origin, pain arising after LC is multifactorial and mainly visceral in nature. Surgical incisions, CO₂ insufflation and raised pressure in abdominal cavity are some of the factors responsible for pain.^{4,5}

Local anesthetics have good analgesic effects and are quite safe to use. Bupivacaine has a half-life of 2.7 – 3.5 hours, 100 mg of bupivacaine in a dose of 2.5 mg/kg of body weight can be given safely in a 70 kg person.⁶⁻⁹ Controversies exist regarding the use of intraperitoneal local anesthetics. Some studies done in this regard showed a significant pain relief post operatively² but some showed little or no use of such agents.¹⁰⁻¹² The objective of this study was to assess the efficacy of bupivacaine in alleviating post-operative pain following LC.

METHODOLOGY

After getting approval of Institutional Review Board (IRB) vide letter No. PF/NA/ADM-I, this randomized controlled trial was conducted in surgical department of Fauji Foundation Hospital, Rawalpindi from July 2017 to Jan 2018. A written informed consent was signed by all patients. WHO sample size calculator was used for sample size calculation by keeping significance level 5%, power of test 80%, anticipated proportions of two groups were, 25.8% and 77.4% respectively.² The sample size came out to be 120 patients divided into two groups ($n = 60$ each). Respondents were selected through non-probability consecutive sampling technique.

Patients with age range 20 – 60 years, both genders and patients according to ASA I and ASA II undergoing LC were enrolled in the study. Those with acute cholecystitis, acute pancreatitis, previous history of abdominal surgery, with intraperitoneal biliary leakage and those allergic to local anesthetic were excluded from the study. They were randomly assigned into two groups, Bupivacaine (group-A) and Control (group-B) by the lottery method (Fig. 1).

Patients in group A underwent LC with administration of 20 ml of 0.5% bupivacaine intraperitoneally. Patients




Fig. 1:

in group B underwent LC without administration of intraperitoneal local anesthetic. Both groups were operated upon by the same team of surgeons using four ports. Post-operative pain was assessed and recorded using visual analogue scale (VAS) at 8 hours postoperatively. A score ≤ 4 at 8 hours were considered as effectiveness. For post-operative pain 30 mg of ketorolac was given I/V and additional doses were given if required.

Statistical Analysis: SPSS version 20 was used for data analysis. For quantitative variables like age and post-operative pain at 8 hours means \pm SD were calculated. Kolmogorov–Smirnov and Shapiro–Wilk test was used for normality data between the groups. Chi square test of statistical significance was used to compare the effectiveness in both groups keeping $p \leq 0.05$ as significant. Effectiveness in both groups was also

stratified among age, gender, BMI and ASA to see the modified effect. Post stratification chi-square test was applied.

RESULTS

A total of 120 patients were evaluated for post-operative pain following laparoscopic cholecystectomy. The socio-demographic characteristics of patients were recorded Table 1.

In group A (IP 0.5% Bupivacaine), mean VAS 8 hours after surgery was 3.63 ± 1.76 and in group B (control) mean VAS 8 hours after surgery was 5.27 ± 1.75 (Table-2). Effective pain control (VAS ≤ 4) at 8 hours after surgery was achieved in 70.0% (n = 42/60) patients in group A as compared to 20% (n = 12/60) in group B (control) (Table-3).

Table1: Socio-demographic details of patients (n = 120).

| Variables | | Groups | | Total |
|--------------------------|------------------------|----------------------|------------------|-------------|
| | | Bupivacaine (n = 60) | Control (n = 60) | |
| Age (Mean) | | 46.80 ± 7.12 | 49.40 ± 6.19 | |
| Gender (%) | Males | 2 (3.3%) | 6 (10%) | 8 (6.7%) |
| | Females | 58 (96.7%) | 54 (90%) | 112 (93.3%) |
| BMI (Kg/m ²) | ≤ 30 Kg/m ² | 48 (80%) | 40 (66.7%) | 88 (73.3%) |
| | ≥ 30 Kg/m ² | 12 (20%) | 20 (33.3%) | 32 (26.7%) |
| ASA Classification | I | 52 (86.7%) | 52 (86.7%) | 104 (86.7%) |
| | II | 8 (13.3%) | 8 (13.3%) | 16 (13.3%) |

Table 2: Mean VAS 8 hours post-surgery and efficacy of treatment (n = 120).

| Bupivacaine Group | Control Group | χ^2 -value | P-value |
|-------------------|---------------|-----------------|---------|
| Mean ± SD | Mean ± SD | | |
| 3.63 ± 1.76 | 5.27 ± 1.75 | 120.0 | 0.001 |

Table 3: Efficacy of treatment in both groups at 8 hours post-surgery (n = 120).

| Efficacy | Bupivacaine Group | Control Group | Total | χ^2 -value | P-value |
|------------------|-------------------|---------------|------------|-----------------|---------|
| Absence of pain | 42 (70.0%) | 12 (20.0%) | 54 (45.0%) | 30.31 | 0.001 |
| Presence of pain | 18 (30.0%) | 48 (80.0%) | 66 (55.0%) | | |
| Total | 60 (100%) | 60 (100%) | 120 (100%) | | |

DISCUSSION

Laparoscopic cholecystectomy is one of the most common surgeries with minimal tissue injury, rapid postoperative recovery, early mobilization, low postoperative complications, and shorter hospital stays. Our results showed demographic characteristics were similar in both groups ($p \geq 0.05$) and effective pain control was achieved in 70% patients in group A as compared to 20% in group B (control) at 8 hours after surgery. The difference was found to be statistically significant ($p = 0.001$).

Similar trends were found when our results were compared with the already published reports on the given subject. Garza et al also found a significant difference ($p = 0.018$) in pain levels between both groups 6 hours postoperatively. Average analgesic requirement was also lower in the bupivacaine group.¹³

In double blind randomized controlled trial Khatri et al observed that intraperitoneal bupivacaine significantly reduces pain following LC. Mean intensity of pain on VAS over 10 hours was higher in placebo group compared to bupivacaine group. All patients in placebo group required analgesia within 6 hours after surgery and 68% in bupivacaine group required analgesia during the same post op period.¹⁴

Gupta et al divided patients into 3 groups as, Group 1 was control, Group 2 was assigned to receive portside infiltration of bupivacaine, while group 3 received combined port site and intraperitoneal instillation of bupivacaine. At 1st, 4th & 8th post-operative hour, minimum VAS score was in group 3. Patients in group 3 received a lower total amount of rescue analgesia and they also had the shortest hospital stay after LC, compared to the patients in the other groups.¹⁵

Shran et al reported VAS was significantly lower after bupivacaine and rescue analgesia was given when VAS was ≥ 6 .¹ Lin et al aimed to compare the efficacy of 50 mg 0.5% bupivacaine in patients undergoing LC for postoperative pain relief. The postoperative pain at intervals of 1, 2, 4, 8, 12, and 24 h showed that VAS pain scores were significantly lower in Bupivacaine group as compared to control group at 1, 2, and 4 hours after surgery with for all variables.¹⁶

Our sample size was relatively smaller and we did consecutive sampling due to time constraint for completion of study in the required time frame, which may result in introduction of bias. We recommend continuation of this project for gathering more patients to increase the sample size to a statistically significant figure. We also recommend random sampling to reduce the bias and to generalize our results to the population in question.

CONCLUSION

Present study showed that intraperitoneal administration of 0.5% Bupivacaine was an effective treatment option for postoperative pain control at 8 hours after laparoscopic cholecystectomy.

Author Contributions:

Conception and design: Nadia Attiq.

Collection and assembly of data: Nadia Attiq, Muhammad Imran.

Analysis and interpretation of data: Attique Sadique, Khalil-ur-Rehman, Naveed Arshad.

Drafting of the article: Aurangzeb Khan, Khalil-ur-Rehman, Naveed Arshad.

Critical revision of article for important intellectual content: Aurangzeb Khan.

Statistical expertise: Attique Sadique, Muhammad Imran, Naveed Arshad.

Final approval and guarantor of the article: Khalil-ur-Rehman, Naveed Arshad.

Corresponding author email: Naveed Arshad: oliajann@gmail.com

Conflict of Interest: None declared.

Rec. Date: Jun 1, 2021 Revision Rec. Date: Sep 25, 2021 Accept Date: October 14, 2021.

REFERENCES

- Sharan R, Singh M, Kataria AP, Jyoti K, Jarewal V, Kadian R. Intraperitoneal Instillation of Bupivacaine and Ropivacaine for Postoperative Analgesia in Laparoscopic Cholecystectomy. *Anesth Essays Res.* 2018; 12: 377-380.
- O'Connell PR. Principles of laparoscopic and robotic surgery. In: Williams, N., O'Connell, P. McCaskie A, editors. *Bailey & Love's short practice of surgery.* 27th ed. UK: Taylor and Francis Group, 2017: p105-118.
- Sinha RY, Raje SR, Rao GA. Three-dimensional laparoscopy: principles and practice. *J Minim Access Surg.* 2017; 13: 165-169.
- Shamim B, Khan AA, Saleem RM. Effectiveness of intraperitoneal administration of 0.5% bupivacaine in post-operative analgesia after laparoscopic cholecystectomy. *Pak Armed Forces Med J.* 2016; 66 (suppl-3): 248-251.
- Jain S, Nazir N, Singh S, Sharma S. A prospective randomised controlled study for evaluation of high-volume low-concentration intraperitoneal bupivacaine for post-laparoscopic cholecystectomy analgesia. *Indian J Anaesth.* 2018; 62: 109-114.
- Nupur C, Shubha S, Vinayak SR. Evaluation of intraperitoneal bupivacaine for postoperative analgesia in patients undergoing laparoscopic cholecystectomy: a prospective randomized trial. *Anaesth Pain Intensive Care.* 2019; 18: 361-366.
- Toleska M, Kartalov A, Kuzmanovska B, Panovski M, Shosholcheva M, Dimitrovski A, et al. Efficacy of intraperitoneal bupivacaine on pain relief after laparoscopic cholecystectomy. *Contributions Sec Med Sci.* 2018; 39: 123-129.
- Upadya M, Pushpavathi SH, Seetharam KR. Comparison of intraperitoneal bupivacaine and intravenous paracetamol for post-operative pain relief after LC. *Anaesth Essays Res.* 2015; 9: 39-43.
- Mostafa RH, Mekki YM. Comparative evaluation of intraperitoneal bupivacaine and bupivacaine ketamine combined with lung recruitment for reducing postoperative shoulder pain in laparoscopic cholecystectomy. *Egypt J Anaesth.* 2018; 34: 159-164.
- Banoria NK, Prakash S, Sharma OK. Effect of local anesthetic instillation intraperitoneally on post-op pain relief after laparoscopic cholecystectomy. *Indian J Clin Anat Physiol.* 2016; 3: 322-325.
- Das NT, Deshpande C. Effects of intraperitoneal local anaesthetics bupivacaine and ropivacaine versus placebo on postoperative pain after laparoscopic cholecystectomy: a randomised double blind study. *J Clin Diagn Res.* 2017; 11: 08-12.
- Ahmad J, Khan ZA, Khan A. Efficacy of port-site and intraperitoneal application of bupivacaine in reducing early post-laparoscopic cholecystectomy pain. *Pak Armed Forces Med J.* 2015; 65: 211-215.
- Garza GC, Diaz-Elizondo JA, Cuello-Garcia CA, Cabello OV. Irrigation with Bupivacaine at the Surgical Bed for Postoperative Pain Relief after Laparoscopic Cholecystectomy. *JSLs.* 2012; 16: 105-111.
- Khatrri AM, Dawani AP, Gazder DP, Jaffery AH, Wasty WH. Intraperitoneal bupivacaine instillation significantly reduces pain following laparoscopic cholecystectomy. *Pak J Med Dent.* 2019; 7: 26-31.
- Gupta R, Rai A, Vasuniya V, Choudhary D. Comparative evaluation of subcutaneous infiltration (port site) vs intraperitoneal infiltration of bupivacaine on post laparoscopic cholecystectomy pain. *Int J Surg.* 2018; 5: 500-506.
- Lin S, Hua J, Xu B, Yang T, He Z, Xu C, et al. Comparison of bupivacaine and parecoxib for postoperative pain relief after laparoscopic cholecystectomy: a randomized controlled trial. *Int J Clin Exp Med.* 2015; 8: 13824-13829.